SEARCHING FOR DRUG BENEFIT BENCHMARKS – COST PER DAY OF THERAPY

The concept of a therapeutic maximum allowable cost, or “therapeutic MAC,” was revived in late 2001 in the form of a “best-in-class” list of drugs in the Michigan Medicaid program and a “referenced-based formulary” method in the Oregon Medicaid program. Drug therapeutic MAC was discussed more than a decade ago and pertained to private drug benefit plans in which the member would pay the cost difference between the reference price and the actual price for the drug, with the obvious intent of influencing drug utilization to lower-cost therapeutic alternatives while making drug benefit costs relatively predictable for the drug plan sponsor. For several reasons, this drug benefit design method did not flourish, notably due to the unknown member cost-share amount for any given drug until determined at the point-of-care.

The current version of therapeutic MAC would ostensibly bring the cost difference between the reference price and the actual drug price from the manufacturer in the form of rebates. By either additional member cost-share or additional manufacturer rebates, the reference price is the cost per day of drug therapy. Yokoyama et al. in this issue of *JMCP* report the results of an intervention with physicians designed to reduce the average cost per day of therapy with selective serotonin receptor inhibitors (SSRIs) by encouraging the use of SSRIs other than fluoxetine. Based upon actual drug costs expressed in 1998 dollars, the intervention program was associated with a 7.4% reduction in the average cost per day of SSRI therapy between 1998 and 1999 and an additional 4.0% reduction between 1999 and 2000, before consideration of drug manufacturer rebates. When compared to national average SSRI market share data, the erosion in fluoxetine in this Texas health plan produced net savings estimated to be \$57,191 in 1999, the first year of the intervention program and \$149,453 in 2000, or a combined two-year savings of \$206,644 or \$0.61 per health plan member per year (PMPY) over the two-year period. The authors did not report the administrative costs of the intervention program, precluding an estimate of the return on investment.

The results reported in this study only suggest and do not prove that intervention with physicians in the form of prescriber profiles or “report cards” combined with counter-detailing in the form of education about relative price differences among therapeutic alternatives may influence the average cost per day of drug therapy. This study addresses a key and recurring subject in managed care, the effect of interventions with physicians designed to improve cost outcomes. The authors did not measure outcomes in clinical quality or service-satisfaction. An article published last month in *JAMA* found similar primary and secondary clinical outcomes in a trial of 573 depressed patients randomized to either sertraline, paroxetine or fluoxetine.

Nevertheless, the article by Yokoyama et al. is remarkable in

a number of ways, in addition to providing some evidence that the intervention was associated with a favorable cost outcome. First, this article raises the bar for reports of managed care interventions published in *JMCP* by reporting multiple measures of utilization, thereby providing more information for readers and more assistance in interpretation of the results and conclusions. The authors reported utilization as measured by the number of patients as well as by the number of prescriptions and days of therapy. The authors also reported real-world drug cost rather than some artificial cost derived from Average Wholesale Price or other unreliable method. (Readers may note that the authors used a conservative, not drug- or even SSRI-specific estimate of inflation, which may have underestimated cost savings by underestimating 1999 and 2000 average cost per day in 1998 dollars.) While drug manufacturer rebates influence net health plan drug costs, this study provides benchmark drug cost PMPM data for SSRI antidepressants that may be of interest to other MCOs and managed care pharmacists.

Reporting utilization by multiple measures helps readers to judge the validity of the authors’ conclusions and equally important, the applicability of this study and its findings to other MCOs. Days of therapy and number of patients for each drug are important measures for other managed care pharmacists who attempt to apply these findings to their MCO environment which varies from the Texas health plan in the use of a mail-order pharmacy option, for example. While mail order prescriptions can be converted to community-pharmacy equivalent prescriptions, days of therapy provides a more reliable measure of utilization, at least for antidepressants in general and SSRIs in particular. The subject of the mail-order option in this health plan is not addressed by the authors, making the report of days of therapy for each of the SSRI drugs that much more important in this article.

What about limitations of the study? Comparison of the utilization of SSRIs in this health plan to a national average benchmark is important, but the benchmark employed by the authors may not be the most reliable. Scott-Levin data for November 1998 showed a fluoxetine Rx share of 34.3% v. 42.3% for their chosen benchmark in 1998, and the Scott-Levin data for November 1999 showed a fluoxetine Rx share of 29.8% v. their benchmark of 35.5%. For the 12-month period from 1998 to 1999, the health plan fluoxetine Rx market share relative erosion was 24.2% v. 16.1% relative erosion for their benchmark v. 13.1% for the Scott-Levin Rx share data. This benchmark analysis suggests that the authors may have underestimated the relative difference in fluoxetine erosion for their health plan versus other health plans and thereby underestimated the effect of their intervention program.

This Texas health plan had SSRI utilization that was atypical. Fluoxetine market share as measured by the number of Rxs, days and patients was 42.2%, 41.6% and 37.4% in 1998, higher than the Scott-Levin fluoxetine Rx share of 34.3% for the SSRI drugs. Sertraline was over-represented at 48.9% com-

pared to 30.3% for the Scott-Levin national data. Paroxetine utilization in this health plan was even more disparate at 8.3% of prescriptions in 1998 compared to 28.9% for the Scott-Levin data. An alternate explanation for the data results could be the promotion of sertraline and pill-splitting rather than “counter-detailing” of fluoxetine.

Aside from the benchmark comparisons, Yokoyama et al. could have addressed more thoroughly the subject of physician “report cards” in particular and managed care interventions in general. There is quite a bit of literature on physician report cards used as a means to influence physician behavior, improve quality of care or reduce the cost of care. A study reported last year in *Lancet* involving reminder messages to general medicine physicians on the appropriate use of lumbar and knee x-ray procedures was found to be associated with a 20% reduction in utilization of lumbar and knee x-rays per 1,000 patients per year among 247 general medicine practices that used six radiology departments. The combination of educational reminder messages with audit data (utilization of x-rays by physician, similar to the intervention employed in the Texas health plan) and feedback to physicians did not appear to have an incremental effect on reducing the utilization of apparent unnecessary x-ray procedures. Unlike the study by Yokoyama et al., the study of x-ray procedures among general practice physicians examined the medical appropriateness of the procedures, finding that only 22% to 44% of the x-ray orders were judged to be definitely indicated, according to guideline criteria.

Also left to future authors in *JMCP* is the important subject of the validity and reliability of physician report cards. The authors of the SSRI study in this issue of *JMCP* did not address this subject. Researchers associated with the Agency for Health Care Policy and Research (AHCPR), now the Agency for Healthcare Research and Quality (AHRQ), found that physician report cards for diabetes care used at three separate MCOs were able to account for only four (4) percent of the variance in patient hospitalizations, office visits, laboratory tests and blood glucose level control rates. In the AHCPR study, as in many MCOs, the small number of patients or the small number of observations (data points) can result in detection of random variation more than systematic variation among physicians. This is and will continue to be a significant and real concern for interventions in all managed care organizations. Future authors of articles in *JMCP* on the subject of managed care interventions are encouraged to fully describe their methods to ensure validity and reliability in their physician report cards and to provide assurances that their interventions are able to detect systematic variation and not simply random variation in apparent practice differences among physicians.

Frederic R. Curtiss, Ph. D., R.Ph., CEBS, Editor-in-Chief

References

1. Caffrey A. Michigan is poised to reduce drug cost by setting restrictions on medications list. *Wall Street J* 2001 Nov 12: B8.

2. Gold R. Oregon's new tack on drug prices. *Wall Street J* 2001 Aug 1:B10.

3. Kroenke K, West SL, Swindle R, et al. Similar effectiveness of paroxetine, fluoxetine, and sertraline in primary care. *JAMA* 2001 Dec 19(286): 2947-55

4. Eccles M, Steen N, Grimshaw J, et al. Effect of audit and feedback, and reminder messages on primary care radiology referral: a randomized trial. *Lancet* 2001; 357: 1406-9.

5. Hofer TP, Hayward RA, Greenfield S, et al. The unreliability of individual physician “report cards” for assessing the cost and quality of care of a chronic disease. *JAMA* 1999 Jun 9 (281): 2098-2105.

FACTORS AFFECTING PHARMACIST CONSULTATION SERVICES IN A UNIVERSITY HEALTH INSURANCE PLAN

The whole of science is nothing more than a refinement of everyday thinking. —Albert Einstein

Nothing pains people more than having to think. —Martin Luther King Jr.

If everyone is thinking alike then somebody isn't thinking. —George Patton

Being fully cognizant of a potential oxymoron, there is a joke among statisticians:

A fellow was out duck hunting when a mallard flew by. The hunter fired a shot, but the pattern was about five feet behind the swift bird. Swinging the gun faster, he fired again. Unfortunately, the pattern was about five feet ahead of the bird, but on the average he had a dead duck.

Generally *JMCP* seeks to publish articles with primary data collection less than five years old. “Currency” is valued. Historically, *JMCP* has usually not published an article in the domain of heavy statistical methods, favoring, instead, to publish articles that most managed care pharmacists could incorporate into their practices, both clinical and “business.” Most executives in the pharmacy benefit field, for example, do not make decisions based on *p* values. In this issue of we broke the rule.

In this issue, Schommer has presented us with a thought-provoking article on patient uncertainty regarding prescription medications. At first pass, I was afraid of entering the statistical swamp. The logistic regression analyses had variables without hierarchical correlation, numerous non-significant odds ratios, etc. Primary data collection took place in 1994. However, the issues raised in pharmacy counseling and patient behaviors (such as compliance) are valid and have “currency.”

In statistics a positive finding is inevitably accompanied by a negative finding. Heads also means no tails on the coin flip. In practice, it's not quite so simple. If ibuprofen relieves your aches and pains, not taking the drug does not CAUSE the symptoms.

Simple descriptive statistics, therefore, may not convey the full meaning. Even so, a difference has to be distinguished between “an association with” and “causality.”¹ Regression analyses allow the former, clinical studies the latter. Regression analysis is probably the only way to tease out the factors explaining some patient behavior. In this study, a different design up front, the use of asymmetric scales rather than yes/no

responses, could have stratified answers and allowed for more common statistical comparisons without “regressing the mean”. Additionally, the use of a binary response (Y/N) question always predicts a more likely positive response, rather than a compound question with two parts. For example, “I desire more information” versus “I desire more information and it has to be written.”

As discussed in the article by Schommer, the population sampling technique was disproportionate, resulting in a study population that was missing the age extremes. This is not our world. The problems with appropriate medication use and the elderly have been well described, as have the problems in the proper use of suspensions and inhalers in the pediatric population. Less well described is the 18-to-23-year-old age bracket in which there is often a transition of health care coverage from parent to adult child. The use of the 18 to 29 year olds as the reference group for the regression analyses may have concealed some medication problems. Other noteworthy observations regarding this article include: lack of a prescription cost indicator to determine if costly medications stimulate more information requests, the usual problem of patient self-reported data, but correlation with actual claims data from the plan should have been available (perhaps with appropriate patient consent).

Is the glass half full or half empty? While there are positive results in this article, the value of Schommer's article is in the negative, the devil-in-the-details. Who does not benefit or need pharmaceutical care intervention? Not only must we manage health care delivery but also the recipients of the care. The descriptive data suggest that most patients do not want or desire consultative services (the overwhelming negative responses to “talked with pharmacist,” “asked the pharmacist a question,” and “desired more information” are extremely sobering for our profession). Indeed in the regression models, the minority of the predictor variables, and sometimes, just one, in the consultative component cells reach significance. Two of the prescriptive states (PS2 and PS6) never reach significance in any cell. Clearly a mismatch of perception and reality is occurring. It is important not to waste human and time resources, when one cannot affect an outcome.

The article by Schommer provides an important historical perspective on the practice of pharmacy, as well as insights into patient behavior. There is little reason to believe that either practice or behavior has changed markedly.² While we can statistically nit-pick, remember the Einstein quote. Does anybody really believe these results are not true? In the past few years, various environmental changes have occurred; prescriptive authority, pharmacy practice acts, pharmaceutical care models in retail or clinics, Internet, etc. that have created potential to “progress the mean” of our practice and create informed consumers. But, have we? By a shotgun approach to patient education, can we? The evidence would suggest not. Dr. Schommer is on the right track here, and this article provides useful information for future research.

Alan H. Heaton, Pharm.D., St. Paul, Minnesota

Editor's Note: Dr. Heaton is vice president of clinical marketing and research for Prime Therapeutics, Inc., based in Eagan, Minnesota, and is a member of JMCP's Editorial Advisory Board. Additionally, he is clinical assistant professor in the College of Pharmacy at the University of Minnesota. While he and Dr. Schommer are colleagues, they did not collaborate or otherwise discuss the research conducted and reported by Dr. Schommer in this issue of JMCP.

References

1. Westfall G, Littlefield R, Heaton A, and Martin S. Methodology for Identifying Patients at High Risk for Osteoporotic Fracture. *Clin Ther.* 2001; 23: 1570-88.
2. Heaton AH, Martin SL, Brelje TV, and Ryu S. Potential Drug Interactions in Patients with Onychomycosis: A Retrospective Database Analysis. *American Society of Health-System Pharmacists.* Reno, NV 6-8-99.

COMMUNITY PHARMACISTS AND DIABETES HEALTH CARE

It is frustrating to try to solve a 100-piece puzzle when one has only 35 of the pieces. Getting diabetes patients to follow the American Diabetes Association's Standards of Diabetes Care is an example of a puzzle that needs more of the pieces to fall into place before care will be improved significantly. The Diabetes Health Care System faces many problems and whenever outcome studies are performed, the results indicate that diabetes is the Rodney Daingerfield of chronic conditions since the subject receives such little respect. Diabetes care is simply a national disgrace. Studies referenced in the article by Drs. Nau and Ponte in this issue of *JMCP* and other studies inform us that less than 5% of diabetes patients achieve the level of care that would slow or prevent the complications of diabetes and 94% of patients do not get their feet examined yearly; over half do not have even one HbA1c level performed; less than a third of patients have a dilated pupil eye exam; less than 25% take an aspirin tablet daily; many patients do not self-monitor blood glucose at all and those that do test do not test often enough. A recent study showed that elderly nursing home patients needed better review of their diabetes care¹. More than 95% of diabetes patients are cared for by primary care physicians yet a recent review² of diabetes care provided by these physicians showed a wide variability in the adherence to ADA standards of care, and the study showed that less than 30% of patients met their LDL-C objective of less than 100 mg/dL. Sobering information.

The Standards of Diabetes Care have been published in numerous medical journals for over 7 years. We have proven and well-described care standards that, if attained, diminish the development of diabetes complications. Yet, few diabetes patients follow these standards and most physician medical records do not document that the standards are being followed, let alone achieved. This chasm in care needs versus care delivery could easily lead to malpractice lawsuits in diabetes care.

Enter the pharmacist. The pharmacist is in the right place at the right time to make a difference in the care of diabetes patients. Pharmacists can help **identify** patients at risk of developing diabetes, and the patient is then screened. The need for better need recognition is suggested by work that shows

that in one-third of elderly people with diabetes, the condition is not diagnosed³. Pharmacists can **assess** the care of the patient and develop a treatment plan to improve the outcomes of care. Pharmacists can also **educate** the patient about how to take charge of the diabetes condition and how to use the many diabetes care products that are available. When needed, the pharmacist can **refer** the patient to other health care providers. The barriers to pharmacists working effectively in disease state management are many and include inadequate reimbursement for the time required. A shortage of pharmacists pushes many pharmacists who would otherwise work in diabetes disease management into dispensing tasks rather than clinical care of the patient. Managed care organizations may fail to recognize the need for diabetes care management provided by community pharmacists. The conduct and publication of good studies that demonstrate the value of community pharmacist intervention can make a difference in crossing this care-chasm.

Drs. Nau and Ponte in their article in this issue add one more piece to the total puzzle relating to diabetes care and the impact of pharmacist's interventions. Even though the number of patients enrolled was small and a control group was not included, the study results suggest that pharmacists can make a difference in the care of diabetes patients. This result is all the

more amazing since the pharmacists in this study met with the patient only three times, and each meeting was less than an hour in duration. The pharmacist provided basic diabetes education to the patient and also performed assessments of blood pressure and lab values. Still, some clinical outcomes showed improvement. The article by Nau and Ponte suggest that pharmacists can affect favorably the quality of care of diabetes patients, in this case, health plan patients in a managed care organization. One more piece of the pharmacist diabetes intervention puzzle found.

R. Keith Campbell, R.Ph., FASHP, FAPhA, CDE,
Washington State University College of Pharmacy Pullman, WA

References

1. Guo JJ, Gibson Jt, Barker KN. Prescription drug utilization and associated factors for nursing home patients with diabetes mellitus. *Consult Pharm* 2001; 16: 1068-76.
2. Stamm PL, Stowers AD, Hornsby LB, Murphey LM. Comparison of diabetic patient care with practice standards: a family practice residency program analysis. *Diabetes Trends* 2001; 13 (Suppl B): 16-25.
3. Franse LV, Bari MD, Shorr RI, Resnick HE, VanEijk JTM, Bauer DC, Newman AB, Pahor M. Type 2 diabetes in older well-functioning people: who is undiagnosed? *Diabetes Care* 2001; 24: 2065-70.