Measuring Adherence and Persistence in Drug Therapy

In this issue of *JMCP*, the topic of medication adherence is addressed in an article by White et al. that compares differences in prescription refill patterns of antihyperlipidemic agents between patient groups receiving medications via mail service or retail pharmacies.¹ While the objective of this naturalistic study was to compare refill patterns between two different types of prescription services utilized by plan members, the results and discussion from this article highlight an important measurement that may directly impact the clinical and economic outcomes related to the use of medications for treating chronic diseases.

Measures of medication adherence, also termed “compliance,” add an important dimension to outcomes studies by providing decision-makers with a valuable metric for comparing the utilization patterns of selected therapeutic agents. When medications are prescribed rationally and patients follow their regimens appropriately, the likelihood of achieving positive patient outcomes increases dramatically. However, when patients don’t adhere to their regimens, which may occur for numerous reasons, the opportunity for achieving positive health outcomes may be jeopardized. Measuring levels of adherence to medication therapies can help to explain or predict which plan design, educational intervention, or particular drug product has the potential to provide optimal patient outcomes within an organized health care setting.

In the article in this issue, the authors included patients in the study who had been newly started on an HMG-CoA reductase inhibitor based on a review of pharmacy claims data from a California HMO. Claims data were analyzed for each patient for a period of one year after initiating the therapy regimen. After controlling for some demographic differences between the study groups, the authors found that patients utilizing the mail service had significantly higher rates of adherence than patients using community pharmacies.

The use of pharmacy claims data to measure patient adherence has gained widespread use within pharmacy benefit programs. Large prescription claims databases have provided researchers with convenient access to utilization patterns of large patient populations and the opportunity to investigate medication adherence.

Perhaps the most comprehensive methodological work in the area of measuring medication compliance using administrative databases comes from Steiner et al. in two widely cited articles.^{2,3} In an original research article and in a subsequent review piece, Steiner and his colleagues provided a rationale for using different methods to calculate the patient adherence rate. Different rates may be calculated depending upon the length of time that one chooses to observe adherence within a population.

Utilizing a measurement based on Steiner’s earlier work, Sclar et al. described the now often-used “Medication Possession Ratio” (MPR) to calculate days of therapy between refills using prescrip-

tion claims histories.⁴ To calculate this ratio, the total days supply of the medication dispensed (not including the last prescription dispensed) is divided by the total number of days between the first prescription dispensed and the last prescription dispensed during an observation period. Typically, this calculated ratio is less than 1.0, owing to the fact that patients will exhibit gaps in medication days. However, if patients obtain refills before their supply has been exhausted, the calculation may exceed 1.0.

As a benefit to readers, White et al. chose to calculate rates of adherence in more than one way. For example, the authors calculated a mean MPR of 93% for patients using mail-service pharmacy and an MPR of 82% for community pharmacy patients. However, over the one-year follow-up period, mail-service patients refilled prescriptions with an average total days supply that covered only 81% of the observed days, while community pharmacy patients’ refills covered, on average, only 57% of the observed days. While mail-service patients had longer mean days of persistence compared to community pharmacy patients (280 days versus 214 days, respectively), the authors did not report rates of discontinuation within the study groups, which would have allowed for an insightful comparison. Previous research has shown discontinuation rates of 10% to 60% within this class of medications.^{5,6,7,8}

Relying solely on a single ratio of adherence, such as an MPR calculation, provides the decision-maker with essentially one dimension of information regarding appropriate and adequate use of a medication within a population. When measuring medication adherence, we have to be sensitive to the different dimensions that should be considered. The first dimension is one of “consistency” with the prescribed regimen. Is the patient actually taking the drug in the manner that it was prescribed? To measure this dimension, the MPR can be used to compare what we would expect utilization to be based on the submitted days supply found in the prescription claim record. The second dimension is one of “persistence” in taking the medication. Is the patient refilling the medication in a timely fashion, over a pre-defined period of time, without gaps in treatment days? Both of these dimensions are equally important in comparative evaluations of medication adherence.

Consider a patient who refills two consecutive 30-day prescriptions in a timely manner, but then discontinues the medication, without the physician’s consent, for the rest of the year. Next, consider another patient that uses the medication intermittently for a total of nine months during the year, creating gaps in treatment. The latter patient’s MPR would be significantly lower than the former patient’s calculated MPR. Of the two patients, which one is more “adherent”? Which patient would be considered more “persistent”? Is one attribute of greater importance in determining appropriate use? Further consider which dimension of adherence would affect long-term total and component health care costs for the patient.

In a recent review of the many methods used to measure medication adherence among patients, Farmer described the advan-

tages and disadvantages of techniques such as biological markers, patient diaries, pill counts, electronic monitoring, and prescription record reviews.⁹ The method chosen to measure adherence can significantly affect the results of the study. Using prescription refills to measure adherence, for example, has some major limitations. First, prescription refills do not provide data on the quantity of medication actually consumed or the timing of doses.¹⁰ Many patients will obtain refills when reminded even when they have medication remaining (especially when they have low cost-sharing requirements); others will stockpile medications or store them in several places for convenience.¹¹ Second, measuring prescription refills from claim records ignores the effect of prescription samples. Third, measuring prescription claim records does not capture information on the prescriptions written but not filled. It has been asserted that about one-third of patients do not have their prescriptions filled.¹² Fourth, one cannot tell from claims data whether premature discontinuance of medication represents a change in physician orders or a decision (rational or irrational) of the patient.

As with any other research, studies on adherence must avoid selection bias. In other words, the groups must be comparable. Comparing groups that differ in days supply dispensed (e.g., 90-day supply for mail-order refills versus a 30-day supply for network prescriptions) is likely to bias the results. A 90-day supply will almost always appear to have a longer duration and greater persistence than a 30-day supply. Whenever possible, confounding variables must be controlled. The authors reported a significantly higher proportion of Medicare+Choice patients in the mail-service cohort (87%) than the community pharmacy cohort (56%), however, this difference was not controlled for in the ANCOVA analyses presented in Table 4. Also, no mention was made as to how differences in copayment levels were controlled. Failure to control for type of insurance coverage (e.g., Medicare+Choice vs. commercial) or copayment level could affect the results considerably; we lose the opportunity to determine if there is, in fact, a true cause-and-effect relationship between the type of prescription service and levels of adherence and persistence.

Measurements must also be valid—they must measure what they purport to measure. If one accepts that “adherence” is a measure of the degree to which patients actually follow prescription orders when self administering medication, then the limitations of measurements such as “duration” and “persistence” become apparent. The appropriateness of the research design and interpretation of the results cannot be over emphasized when assessing rates of adherence and persistence within a managed care population; these measurements have the potential to influence the assessment of the overall clinical and economic benefit of prescription utilization.

As we continue to measure the economic impact of medications and interventions within managed care, the consideration of adherence and persistence rates should be an integral part of the assessment process. It seems reasonable to think that poor adher-

ence rates within patient populations would ultimately lead to greater subsequent medical costs. However, an overview of the relationship between adherence rates and medical expenditures conducted by Hughes et al.¹³ concluded that while poor adherence rates almost always had a negative effect on the efficacy of the drug therapy, the economic impact varied. The results were based on a review of 22 studies previously published in the literature. The most notable scenario where economic impact was least affected by poor adherence was in the case where drug costs dominated total costs within a disease state. In summary, it appears that the economic effects of poor adherence within a population are unpredictable, at best.

Insightful administrators would do well to encourage the inclusion of real-world levels of adherence into the decision-making process when considering both the potential clinical and economic outcomes related to drug therapies or interventions within a managed care patient population. As a follow-up to their earlier research, Hughes et al.¹⁴ have suggested methodologies that can account for estimated levels of adherence within patient populations.¹⁴ This recognition of varying adherence rates allows decision-makers to conduct sensitivity analyses through modeling exercises to better predict outcomes in a naturalistic setting.

In addition, more research is necessary to better determine a range of “acceptable” levels of adherence rates within therapeutic categories. We know that 100% compliance is optimal, but not likely attainable across a population. Therefore, is 90% adherence adequate and attainable? How about 75%? Authors often select an anecdotal level of 80% as an appropriate indicator of compliance, but offer no rationale for the selection of that particular rate. Studies that evaluate the relationship between different rates of adherence and their direct effect on clinical and economic outcomes should be encouraged in order to determine practical levels of acceptable adherence rates.

As these investigations continue, we may learn how to better achieve optimal levels of medication adherence, thus improving the potential for positive patient outcomes within the managed care setting. We must recognize the shortcomings of our present methods of measurement of adherence and persistence in drug therapy management.

Michael Johnsrud, Ph.D.

Associate Director

The Center for Pharmacoeconomic Studies

The University of Texas at Austin

E-mail: johnsrud@mail.utexas.edu

Kenneth W. Schafermeyer, Ph.D.

Professor and Director of Graduate Studies

St. Louis College of Pharmacy

E-mail: Kschafermeyer@sticop.edu

(See references, next page)

REFERENCES

1. White TJ, Chang E, Leslie S, et al. Patient adherence with HMG reductase inhibitor therapy among users of two types of prescription services. *J Man Care Pharm* 2002;8(3):186-91.
2. Steiner JF, Koepsell TD, Fihn SD, et al. A general method of compliance assessment using centralized pharmacy records. *Med Care* 1988;26:814-23.
3. Steiner JF, Prochazka AV. The assessment of refill compliance using pharmacy records: Methods, validity, and applications. *J Clin Epidemiology* 1997;50:105-16.
4. Sclar DA, Skaer TL, Chin A, et al. Utility of a transdermal delivery system for antihypertensive therapy: Part 1 *Am J Man Care* 1991;91:50-56.
5. Avorn J, Monette J, Lacour A, et al. Persistence of use of lipid-lowering medications: a cross-national study. *JAMA* 1998;279:1458-62.
6. Simmons LA, Levis G, Simons J. Apparent discontinuation rates in patients prescribed lipid-lowering drugs. *Med J Aust* 1996;164:208-11.
7. Oster G, Borok GM, Menzin J, et al. Cholesterol-reduction intervention study (CRIS): a randomized trial to assess effectiveness and costs in clinical practice. *Arch Intern Med* 1996;156:731-39.
8. Hiatt JG, Shamsie SG, Schectman G. Discontinuation rates of cholesterol-lowering medications: implications for primary care. *Am J Man Care* 1999;5:437-44.
9. Farmer KC. Methods for measuring and monitoring medication regimen adherence in clinical trials and clinical practice. *Clin Ther* 1999;21(6):1074-90.
10. Claxton AJ, Cramer J and Pierce C. A systematic review of the associations between dose regimens and medication compliance. *Clin Ther* 2001;23(8):1296-1305.
11. Cramer JA, Spiker B, eds. Patient compliance in medical practice and clinical trials. New York: Raven Press; 1991.
12. Lofholm PW and Katzung BG. Rational Prescribing and Prescription Writing. Chapter 66 in Katzung BG (ed.) *Basic & Clinical Pharmacology*. New York: Lang Medical Books, 2001.
13. Hughes DA, Bagust A, Haycox A, et al. The impact of non-compliance on the cost-effectiveness of pharmaceuticals: a review of the literature. *Health Econ* 2001;10:601-15.
14. Hughes DA, Bagust A, Haycox A, et al. Accounting for noncompliance in pharmaco-economic evaluations. *Pharmacoeconomics* 2001;19(12):1185-97.

Conjoint Analysis in Pharmaceutical Research

The Holdford and Carroll article¹ in this issue of *JMCP* uses conjoint analysis, a methodology for evaluating subject preference assessment that is unfamiliar to many pharmacists, other clinicians, and health services researchers. Despite the lag in adoption of these methods in the health care field, conjoint analysis is rapidly gaining popularity as a versatile research tool that can be used in a variety of evaluation contexts. There are a number of important pharmacy and pharmaceutical research questions that naturally suggest conjoint analysis as a valuable research methodology. These include new drug development, drug benefit program design, drug treatment willingness-to-pay assessment, and medication cost effectiveness research. A current Medline search shows that while only 26 articles with the keywords "conjoint analysis" appeared in the medical literature prior to 1997, 40 articles have

appeared from 1997 to date, with the citation rate increasing by roughly 30% annually.

Conjoint analysis methods were originally developed by market researchers over 40 years ago, and have been further advanced and widely used in marketing studies by academic and commercial marketing research groups.² These methods received formal underpinnings in economic utility theory early in their development.^{3,4} Powerful tools for empirical evaluation of conjoint analysis applications were pioneered principally by Daniel McFadden, who shared the 2000 Nobel Prize in economics for this research.^{5,6} The initial application for McFadden and colleagues was evaluation of consumer preferences for alternative transportation options, specifically, the introduction of the BART subway system for northern California commuters.⁷

What Is Conjoint Analysis?

Conjoint analysis is a method for systematically evaluating and estimating the strength of subject (e.g., consumer, patient, clinician, decision-maker, etc.) preferences for discrete choice alternatives (e.g., different therapies, different drugs, health insurance choices, disease symptoms, treatment side-effects, etc.). The key insight behind conjoint analysis is that people systematically chose among alternatives based on observable "bundles" of attributes that each choice option represents.⁸ For example, consumers shopping for a new automobile would consider the vehicle size, body style, color, price, engine power, fuel efficiency, seating capacity, safety features, and other observable characteristics, including the brand name itself (e.g., Ford, BMW, Chevrolet, Honda, etc.) in making a car purchase decision. While each individual possesses idiosyncratic or unobservable tastes that contribute to observed choice, the researcher can systematically evaluate the strengths of each of these unobservable attributes in eliciting choice preferences through a conjoint analysis experiment, where a variety of vehicle profiles are ranked by potential car-buyers on the basis of varying the bundles of observable (and thus modifiable) attribute bundles.

In a conjoint analysis experiment, the potential car buyers would be asked to rank their preferences for vehicles with different combinations of the salient observable characteristics. Subjects with different demographics (e.g., income, family size, education, occupation, etc.) will almost certainly demonstrate different preference strengths for different choice attributes. This allows the car manufacturer market researchers to "segment" the market into groups of consumers that have similar tastes, and then provide a variety of vehicle choices that better match the spectrum of tastes in the consumer population.

Why Has It Taken So Long for Conjoint Analysis to Be Used in Health Care?

Conjoint analysis generally utilizes stated preference (SP) methods, rather than revealed preference (RP) methods. In SP studies, subjects are provided information on alternative choice profiles in

an experimental, hypothetical-choice setting. They are asked to state their choice preferences “as if” they were actually going to be given their choice in the real world. In RP studies, the choices that subjects make in the real world are the units of analysis. For example, the quantity and variety of breakfast cereals people actually buy as a function of prices and other consumer characteristics could be analyzed using retrospective data from grocery stores chains, an RP method. Similarly, the amount of health care that people utilize as a function of health plan copayments and other characteristics can be observed and analyzed in a prospective RP study, such as the Rand Health Insurance Experiment.⁹

Many social science and health services researchers have long held strong biases against SP research methods. They have believed that subjects do not respond in hypothetical experimental settings as they really would if confronted with such choices in the real world. They think that SP subjects can easily see what answer is “best” for them, and then strategically rank their answers to give the researcher a false impression of their true preferences.

For example, subjects in a naive SP experiment on health insurance plan design might systematically overstate their preferences for freedom-of-choice of provider, generosity of coverage, or accessibility of facilities if they were not actually paying for these features. Similarly, taxpayers might systematically understate their preferences for increasing taxes to cover a new hospital, highway, or public works project if they thought that doing so would convince politicians to vote against these programs.

Conjoint analysis can overcome these problems through experimental designs that make it impossible for the SP subject to game their responses by comprehending what the “correct” answer should be. In a well-designed conjoint analysis experiment, the choice profiles vary in such a way that strategic responses aren’t possible.

From another perspective, there are many critical research questions that simply cannot be evaluated with RP methods. For example, suppose a pharmaceutical manufacturer wants to know whether a new drug in early development has efficacy or toxicity characteristics that clinicians and patients would prefer to the existing FDA-approved therapeutic alternatives. Before investing significant resources in clinical trials, the company could sponsor clinician and patient conjoint analyses of the new drug profile compared to the existing alternatives. RP data for this drug would not become available until the drug was actually approved, which would potentially require millions of additional research dollars and years of investment.

One of the key recent advances in conjoint analysis research is the insight that conjoint analysis research studies can be combined with RP data to provide results that are more useful and valid than those of either study design used separately.⁸ Thus, if researchers had a very good RP study on patient preferences for existing antihypertensive drugs, they could use that, in combination with a conjoint analysis of a new experimental antihypertensive medication to predict how well the new therapy would be accepted by

various types of patients, more precisely than using just the RP study or just the conjoint analysis alone.

Validity of Conjoint Analysis

Conjoint analysis studies have been subjected to substantial validation research.⁸ Perhaps the strongest proof of validity of this research method is the fact that nearly all major corporations utilize either internal or external conjoint analysis market research in product design and pricing decisions. Governments and public agencies also rely heavily on conjoint analysis methods for transportation planning, environmental policy planning, and other public policy research. If the conjoint analysis techniques were not valuable, they would not be used routinely for these multimillion dollar public and private decisions.

Conducting a Conjoint Analysis

Designing a successful conjoint analysis experiment involves first establishing the important characteristics of choice that lead people to prefer one alternative to others. This is often done by assembling a focus group of representative subjects (e.g., consumers, patients, or clinicians) and with a structured interview format to elicit the most important attributes-of-choice decisions. The resulting information is then used to create a pilot data collection instrument that is administered to a small sample of subjects to ensure that the survey instrument is capturing meaningful and relevant dimensions of subject choice. A full-scale survey is then implemented using statistical power calculations to estimate necessary sample sizes and fractional factorial experimental designs to ensure that adequate data are collected on each decision attribute of interest.^{8,10} The data are analyzed using multinomial logit or other qualitative dependent variable estimation techniques.¹¹ Policy analyses are then performed to (a) evaluate and rank the strengths of each attribute in determining subject choice, (b) establish how these rankings change for different demographic subgroups, (c) predict how many subjects would switch to another choice (e.g., a new drug) if specific attributes were available for that choice, (d) estimate the strength of the price or cost attribute to determine subject willingness to pay for various choice alternatives, and (e) estimate subject utility directly as a function of choice alternatives.

Use of Conjoint Analysis in Medical, Pharmaceutical and Health Services Research

Holdford and Carroll list several areas where conjoint analysis has been successfully applied in areas of clinical interest, including consumer preferences for drug therapies, pharmaceutical services, health outcomes, and health insurance.¹ Additional subjects found in a Medline search include: health status preference scales,^{12,13} new drug development,¹⁴ patient preferences for clinical products and services,¹⁵⁻¹⁷ and decision-maker preferences.¹⁸

Conjoint analysis has many advantages over other health status assessment and ranking techniques. Like the standard gamble and

time trade-off methods, it is rigorously grounded in economic utility theory.¹⁹ It is much less cumbersome to implement in cost-effectiveness applications and other settings than these other utility measures. In fact, any health status measurement tool can be converted into a conjoint analysis index of consumer preference. It is also rigorously grounded in econometric estimation models with testable parameter restrictions.⁸ Conjoint analysis provides direct assessment of consumer willingness-to-pay for treatment options and consumer utility, in an SP framework that is resistant to subject "response gaming." It can be combined with data from empirical RP studies to provide results that are more precise than either method separately. It has been used and validated in numerous private-sector and public-sector studies in many different fields of preference elicitation. In short, we can expect to see conjoint analyses used increasingly in health care, pharmaceutical, and pharmacy evaluations.

Joel W. Hay, Ph.D.
Dept. of Pharmaceutical Economics and Policy
Univ. of Southern California
E-mail: jhay@usc.edu

REFERENCES

1. Holdford D, Carroll N. Consumer preferences for types of cost containment in prescription drug programs. *J Man Care Pharm* 2002;8(3):192-98.
2. Cattin P, Wittink DR. Commercial use of conjoint analysis: A survey. *J Marketing* 1989;46:44-53.
3. Lancaster KJ. A new approach to consumer theory. *J Polit Economy* 1966;74(2):132-57.
4. Rosen S. Hedonic prices and implicit markets: Product differentiation in pure competition. *J Polit Economy* 1974;82(1):34-55.
5. McFadden D. Conditional Logit Analysis of Qualitative Choice Behavior. In Zarembka, P. (ed.), *Frontiers in Econometrics*, New York: Academic Press 1974;105-42.
6. McFadden D. The choice theory approach to market research. *Marketing Science* 1986;5(4):275-97.
7. Train K, McFadden D. The goods/leisure trade-off and disaggregate work trip mode choice models. *Transportation Research* 1978;12:349-53.
8. Louviere J, Hensher D, Swait J. *Stated Choice Methods*, New York: Cambridge University Press, 2000.
9. Newhouse J, and the Health Insurance Experiment Group. *Free for All? Lessons from the Rand Health Insurance Experiment*. Cambridge MA: Harvard University Press 1993.
10. Gustafsson A. *Conjoint Measurement: Methods and Applications*. New York, NY: Springer, 2000.
11. Maddala GS. *Limited-dependent and qualitative variables in econometrics*. New York, NY: Cambridge University Press, 1983.
12. Fraenkel L, Bodardus S, Wittink DR. Understanding patient preferences for the treatment of lupus nephritis with adaptive conjoint analysis. *Medical Care* 2001;39(11):1203-16.
13. Gotoh F, Terayama Y, Amano T. The Stroke Scale Committee of the Japan Stroke Society. Development of a novel, weighted, quantifiable stroke scale: Japan stroke scale. *Stroke* 2001;32(8):1800-07.
14. Bingham M, Johnson FR, Miller D. Modeling choice behavior for new pharmaceutical products. *Value in Health* 2001;4(1):1-13.
15. Leung GM, Chan SS, Chau PY, Chua SC. Using conjoint analysis to assess patients' preferences when visiting emergency departments in Hong Kong. *Academic Emergency Medicine* 2001;8(9):894-98.
16. Shackley P, Slack R, Michaels J. Vascular patients' preferences for local treatment: an application of conjoint analysis. *J Health Services & Research Policy* 2001;6(3):151-57.
17. Meister H, Lausberg I, Walger M, von Wedel H. Using conjoint analysis to examine the importance of hearing aid attributes. *Ear & Hearing*, 2001; 22(2):142-50.
18. Bala MV, Mauskopf J. Estimating the Bayesian loss function. A conjoint analysis approach. *International J Tech Assess in Health Care*. 2001;17(1):27-37.
19. Torrance G. Measurement of health state utilities for economic appraisal: A review. *J Health Economics* 1986;5(1):1-30.

Better Data for Making Better Decisions: Finger-Pointing or Useful Drug Use Review (DUR)

The research presented by Heaton et al.¹ in this issue of the *Journal* suggests that a significant number of patients were prescribed itraconazole concomitantly with potentially interacting medications. Another study,² using similar retrospective database research methods, drew comparable conclusions about apparent drug interactions with cisapride. The implications of these research reports are several-fold and raise important questions about pharmacy practice in general and drug use review (DUR) specifically.

The first implication is that online prospective drug use review (OPDUR) systems are not systematically identifying specific drug-drug interactions (DDIs). This may be because the alerts generated by OPDUR systems do not have high sensitivity or specificity. Second, the results of this study and others like it, with better data and methodology, could help make prospective DUR screening more effective by focusing on only those problems characterized as clinically significant. Third, many pharmacists do not consider OPDUR to be helpful or clinically relevant to their pharmacy practice in the "real world." A major concern of experts is that this barrage of OPDUR alerts does nothing to advance patient care and may actually desensitize pharmacy personnel to such messages. This has limited the potential of any OPDUR system to improve patient care. Lastly, OPDUR systems have practical and functional limitations that impact their utility. In this review, we explore these issues in brief and set the stage for a more thorough review of state-of-the-art DUR to be published in a future issue of *JMCP*.

Sensitivity and Selectivity of OPDUR Systems

Evidence of significant problems with OPDUR systems and their components has been mounting for some time. The USP (United States Pharmacopeia) Drug Utilization Review Advisory Committee (1995-2000) has pointed to problems with existing DUR criteria.³ The committee concluded that wide variation exists in the current sets of DUR criteria. A leading cause of this inconsistency has been the lack of formal methods governing the development of DUR criteria. Additionally, the group called for a shift

away from an individual patient review DUR to a more comprehensive disease-management approach.³ The concerns of the USP panel are illustrated by a study conducted by Hazlet et al.⁴ They tested the ability of prospective DUR systems in chain and HMO pharmacies in the state of Washington to detect 16 well-established drug-drug interactions (DDIs). These systems failed to detect the specified interactions one-third of the time. The chance of detecting important DDIs (sensitivity) ranged from 0.44 to 0.88. The specificity of these systems (few false positives) ranged from 0.71 to 1.0. The challenge in developing optimal or at least more useful OPDUR systems is to refine the criteria in such a manner that an alert message is highly predictive of an event.

Clinical Significance of OPDUR Alerts

The results of the Heaton et al. study and others like it could, with better data and methodology, help make prospective OPDUR screening more effective by focusing on only those problems characterized as clinically significant. Heaton and colleagues indirectly point to the difficulties associated with assessing the true magnitude of any specific DDI problem. While clinical studies well document the potential severity of DDI problems, our knowledge of the magnitude of such problems in the “real world” is relatively sparse. Experience suggests that many patients exposed to dangerous drug-drug interactions do survive them, often without using medical care services.

A fundamental goal of all DUR systems should be to safeguard patient safety at the point of service without compromising the patient's interest.⁵ The practical task is how to accomplish this within the context of currently available electronic systems. The challenge before us is to answer basic questions, such as, what is the frequency of a specific adverse clinical consequence among patients exposed or receiving alerts (incidence).

Common clinical epidemiological techniques can help us in this respect. For example, in assessing the value of diagnostic tests, we look for measures that have high sensitivity values. Applied to a drug-drug interaction, the sensitivity score would reveal the percent of all clinical cases (e.g., bleeding episodes) that would have been detected by an alert message. An analogous and possibly more intuitively meaningful measure of “true positive” alerts is the positive predictive value. This is the percentage of all alerts that, if left unaddressed, would result in this clinical consequence. For example, of all flagged warfarin-aspirin alerts, what percentage of them would have resulted in a bleeding episode? Similar specificity and negative predictive value measures help define “rule out” criteria (true negatives).

Claims databases can help in making these types of assessments. Either sensitivity or specificity or positive/negative predictive values can and should be used as yardsticks for assessing the utility of existing or more sophisticated screening criteria as called for above. Having said that, we must hastily point out that this type of analysis is not easy to perform.

Pharmacist Response

Pharmacists have criticized OPDUR systems for generating alert messages that they consider to be simply “noise” in the system. Often the prescriber, the patient, and even the pharmacist are unaware of the OPDUR alerts. On-site systems may not contain comprehensive drug profiles if the patient receives prescriptions from several pharmacies or pays cash for other prescriptions. Many alerts focus on clinically insignificant problems about drugs generally regarded by the physician and pharmacist as safe in a given clinical situation.⁵ Not surprisingly, many messages are ignored or these functions are deactivated. In a study of community pharmacy practice in Indiana, Chui and Rupp report that 88% of the alerts were overridden.⁶ The reasons for the overrides included: (1) pharmacist already knew of the problem (34.2%); (2) pharmacist did not think the problem existed (33.6%); or (3) the problem was judged as not clinically significant (27.3%). The challenge in developing optimal, or at least more useful, OPDUR systems is to refine them in such a manner that an alert message is highly predictive of an event.

The workplace environment of pharmacists must also change. Today, dispensing pharmacists work in surroundings where the emphasis is on prescription productivity. Pharmacists have little or no incentive to engage in DUR activities. Moreover, constraints and expectations placed on today's pharmacists preclude thoughtful decisions regarding questionable software-generated OPDUR alerts. When pharmacists do intervene, these activities are often not recorded or tracked in a comprehensive manner. The use of National Council on Prescription Drug Programs (NCPDP) version 3.2 transaction code sets for coding pharmacist interventions and outcomes could help remedy the deficiency in documentation.

Practical Limits

Lastly, OPDUR systems have practical limits. The clinical relevance of OPDUR alerts could be improved greatly if viewed from within the context provided by an electronic patient profile based on comprehensive claims data. However, either many OPDUR systems lack this capability or the necessary patient and prescriber data are missing or inconsistent. Online DUR alert systems, in contrast to in-pharmacy systems, offer a better capability for capturing a complete drug profile for patients. However, the ability to integrate these data sources is limited by demands for data processing efficiency and practical time limits in an online transaction-processing environment.

Lack of sophisticated data processing also limits the clinical relevance of OPDUR alerts. A sophisticated OPDUR system should have the capability to modify its “hits” based on previous pharmacy claims, medical history, laboratory procedures, etc. For example, if a prescriber has already acknowledged his awareness of a particular problem but considers it clinically insignificant, this response can be integrated into the OPDUR system and future alerts suppressed. Further, an ideal system would provide a link to useful information about how to manage a DDI once detected. For

example, a potential drug interaction with warfarin does not carry the same danger if warfarin is the precipitant drug and the prescriber is monitoring INRs. The goal of the sophisticated OPDUR system is to use the most precise selection criteria without sacrificing sensitivity.

Unfortunately, most present-day OPDUR systems are not state of the art and do not live up to the potential of electronic DUR. A quality chasm exists between the operation of OPDUR systems today and the need to protect patients from avoidable clinically significant drug interactions. It is time to expand DUR from individual criteria violations to a systematic review of broadly-based therapy problems. Experts in this area have called for a refocusing of the DUR process.³ Organizations such as the USP and the Academy of Managed Care Pharmacy (AMCP) are working to improve DUR systems. The focus of research should be on imaginative programs to increase the predictive value and selectivity of OPDUR, integration of OPDUR and retrospective DUR, data sharing with the prescriber prospectively, and improving the quality of criteria. Managed care pharmacy programs have the tools to conduct expert DUR. The time has come to expect a higher level of sophistication from our DUR systems so we can assure that patients receive the optimal benefit from their drug therapy.

Final Thoughts

A growing concern exists in the health care community that the medication use system is flawed.⁷ Systems that support effective and appropriate medication use do not exist at the population level; instead, pharmacotherapy is often a fragmented collection of independent actions. The net result is that patients are unnecessarily exposed to known hazards through interacting medications and other drug related problems.

Critics rightly point out the failings of current OPDUR systems or of their use by pharmacists, or both. OPDUR systems can be powerful tools to assure patient drug-related safety. The pharmacist using a comprehensive, sophisticated OPDUR system is the last line of defense for avoiding preventable drug-related morbidity. But these systems are not living up to their potential. We fervently encourage professional organizations and researchers to work collaboratively to develop the clinical and epidemiological basis for new and more sophisticated standards for OPDUR alerts.

Even the most sophisticated OPDUR system is useless if pharmacists do not respond to alerts. Better pharmacist response to alerts requires a combination of more education about improved alert systems, changed performance expectations, and removal of disincentives to respond to alert messages.

Finally, we must recognize that OPDUR systems have limits and cannot be relied upon exclusively to assure patient safety. OPDUR systems often stand alone without a strategic component designed to identify suboptimal prescribing problems or retrospective patterns. Qualitative standards of performance, such as that developed by the Institute for Safe Medication Practices, should be adopted. A strategic system that incorporates quality

assurance elements is the responsibility of, and a challenge for, drug distribution system managers.

*Mark C. Pugh, Pharm.D., R.Ph., Pharmacy Consultant,
Trigon Blue Cross Blue Shield
E-mail: mpugh@trigon.com*

*Dale B. Christensen, Ph.D., R.Ph., Chairman,
Pharmaceutical Policy and Evaluative Sciences,
School of Pharmacy, University of North Carolina
E-mail: dale_christensen@unc.edu*

*Thomas R. Fulda, B.A., M.A., Program Director, DUR,
United States Pharmacopeia
E-mail: trf@usp.org*

*Alan Lyles, ScD, M.P.H., R.Ph., Associate Professor, Health
Systems Management, University of Baltimore
E-mail: alyles@UBmail.UBalt.edu*

REFERENCES

1. Heaton AL, Hansten PD, Martin SL, et al. High frequency of itraconazole prescriptions with potentially interacting medications in a large health care plan. *J Man Care Pharm* 2002;8(3):199-203
2. Jones JK, Fife D, Curkendall S, Goehring E Jr, Guo JJ, Shannon M. Coprescribing and codispensing of cisapride and contraindicated drugs. *JAMA* 2001;286:1607-09.
3. Armstrong EP, Byrns PJ, Foster TS, Morris LS, Fulda TR. Drug utilization review: mechanisms to improve its effectiveness and broaden its scope. The U. S. Pharmacopeia Drug Utilization Review Advisory Panel. *J Am Pharm Assoc.* 2000;40:538-45.
4. Hazlet TK, Lee TA, Hansten PD, Horn JR. Performance of community pharmacy drug interaction software. *J Am Pharm Assoc.* 2001;41:200-04.
5. Soumerai SB, Lipton HL. Computer-based drug-utilization review—risk, benefit, or boondoggle? *N Engl J Med* 1995;332:1641-45.
6. Chui MA, Rupp MT. Evaluation of online prospective DUR programs in community pharmacy practice. *J Man Care Pharm* 2000;6(1):27-32.
7. Hepler CD. Regulation for outcomes as a systems response to the problem of drug-related morbidity. *J Am Pharm Assoc.* 2001;41:108-15.